

Original Article

A narrative review of the control of mosquitoes by Larvivorous fish in Iran and the world

Ali Jafari^{1,2} Ahmadali Enayati³ Fatemeh Jafari¹ Farzad Motevalli Haghi⁴ Nasibeh Hosseini-Vasoukolaei⁴ Reza Sadeghnezhad⁵ Mostafa Azarnoosh¹ **Mahmoud Fazeli-Dinan^{4*}**

1. MSc Student in Medical Entomology, Department of Medical Entomology and Vector Control, Student Research Committee, Faculty of Health, Mazandaran University of Medical Sciences, Sari, Iran

2. Health expert of Administration of Mazandaran Province Prisons, Sari, Iran.

3. Professor of Medical Entomology, Department of Medical Entomology and Vector Control, Health Sciences Research Center, Addiction Institute, Faculty of Health, Mazandaran University of Medical Sciences, Sari, Iran

4. Assistant Professor of Medical Entomology, Department of Medical Entomology and Vector Control, Health Sciences Research Center, Addiction Institute, Faculty of Health, Mazandaran University of Medical Sciences, Sari, Iran

5. Engineer in Environmental Health, Student Research Committee, Faculty of Health, Health Sciences Research Center, Mazandaran university of Medical Sciences, Sari, Iran

*Correspondence to: Mahmoud Fazeli-Dinan
fazelidinan@gmail.com

(Received: 13 Feb. 2019; Revised: 24 Mar. 2019; Accepted: 8 Jun. 2019)

Abstract

Background and purpose: Mosquitoes are responsible for the transmission of many pathogens such as malaria, yellow fever, dengue fever, and so on. The control of mosquitoes using chemical insecticides is not always a sensible approach, so, alternative biological control methods, especially the use of larvivorous fishes, can play a significant role in controlling of mosquito larvae.

Materials and Methods: In this narrative review study, papers and dissertations were collected without time and language limits from international electronic databases in Google Scholar, PubMed, ScienceDirect, Web of Science, Ovid, Medline and WHO site, and Iranian scientific databases including: Barakatkn, SID, Civilica, Magiran, and Medlib using appropriate keywords from 1937 to 2018. Finally, 55 sources were selected and criticized, interpreted, and analyzed.

Results: In the study, some of larvivorous fishes including *Aphanius dispar*, *Carassius auratus* (goldfish), *Gambusia affinis*, and *Poecilia reticulata* (guppy) have been investigated as important predators of mosquito larvae. Among these fish, *Gambusia*, *Aphanius dispar*, *Colisa Lalia*, *Danio rerio*, Goldfish, Guppy and *Oreochromis mossambica* are present in different regions of Iran.

Conclusion: Given the fact that malaria carriers are present in many regions of Iran and the climate of Iran is also potentially suitable for the transmission of malaria, it is recommended to use larvivorous fishes that are compatible with the environmental conditions of each area.

Key words: Mosquitoes; Biological Control; Larvivorous Fishes; Iran

Citation: Jafari A, Enayati AA, Jafari F, Motevalli Haghi F, Hosseini-Vasoukolaei N, Sadeghnezhad R, Azarnoosh M, **Fazeli-Dinan M***. Air born bacteria evaluation in the kitchen air of restaurants in Babol City. Iran J Health Sci. 2019; 7 (2): 49-60. DOI: 10.18502/jhs.v7i2.1064

Copyright © 2019, Published by Mazandaran University of Medical Sciences on behalf of Iranian Journal of Health Sciences and Health Sciences Research Center. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial 4.0 International License <https://creativecommons.org/licenses/by-nc/4.0/> which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

1. Introduction

Yearly more than one billion persons are infected or die from vector-borne diseases, of which mosquito-borne diseases make up a significant proportion (1). So, such diseases transmitted by mosquitoes in (almost) all tropical and subtropical countries are a major problem. Because these species are responsible for the transmission of important diseases including malaria, yellow fever, dengue fever, zika, chikungunya, filariasis, encephalitis, etc. It should be noted that the vector control plans have traditionally focused on killing mosquitoes using various pesticides including organophosphates, carbamates, and pyrethroids (2). Although the application of chemical pesticides is considered as an effective method, because of the adverse effects of chemical pesticides, especially on human health, non-target organisms, increasing the resistance of mosquitoes against these compounds and, consequently, the resumption of transmitted diseases has led to the prohibition of using most insecticides for mosquitoes or at least imposing severe restrictions on many of them by environmental protection agencies (3).

On the other hand, the financial burden of insecticide for the vector control programs is a fundamental constraint to widespread use (2). Therefore, the study and application of sustainable and safe methods, such as biological control is noteworthy in mosquito controlling (4). Biological control methods as a friendly way for the environment can be a suitable alternative to decrease the problems arising from the use of pesticides, such as selection pressure for insecticide resistance (2). Today, the use of biological methods in malaria control programs,

especially in urban and rural areas, has been well-developed in different countries (5), because this method has the potential to be used with other control methods in the mosquito's integrative campaign (6). There is a wide range of living organisms that help regulate the population of mosquitoes through hunting, parasitism, and competition.

Natural organisms that kill mosquitoes have a various range including predators, parasitoids, entomopathogens, etc. (2). In the biological control of mosquitoes using predators, the role of larvivorous fish that consume the larval stage of mosquitoes is very important (2). Larvivorous fishes are one of the most important biological control agents for mosquitos' control, which began to be used widely in the world since the early 1900s and before the use of DDT (7). Many species of fishes have been used for the control of mosquito larvae as a biological control agent and the results indicated that they could be introduced as one of the best candidates for mosquitos' biological control programs (8, 9).

One of the most important species of larval fishes is *Gambusia affinis*, which has a high potential for larvae. This fish, as a native species in the southern regions of the United States, was transferred to the Hawaiian Islands, then to Spain, Italy, and 60 other countries to control malaria (10). Also from early 1908, the species of *Poecilia reticulata*, a native of South America, was introduced so as to control malaria in India, Britain, and many other countries (10). But the use of biological control methods was limited with the introduction of DDT to control malaria in the mid-1940s. Later, eradicating mosquitoes continued with other artificial insecticides, until the first resistances occurred against the insecticides, and

again the use of biological methods was introduced (7). It should be said that natural enemies of mosquitoes, especially in aquatic ecosystems, can play an important role in reducing culicidae populations without making selective pressure (2).

In choosing biological control programs, it is important to select the biological agents. Although larvivorous fish has been demonstrated to be very effective at decreasing mosquito larval density in a variety of habitats (2), in the application of larvivorous fishes, considerations such as searching efficiency, hunting power, adapting to different climatic conditions, and interacting with predators is very important (11). As a result, this method is feasible if there is a proper knowledge of ecology and behavior of the larvivorous fishes and mosquito larvae (12). For example, larvivorous fishes feeding on immature mosquitoes should be small and resistant to drought. Also, they should have ability to live in shallow water among thick grasses, deep water, and various water reservoirs. At the same time, they must be resistant to touch and have the ability to be carried for long distances. They have a high reproduction in their life cycle and can be fed on water, and are interested in being fed from mosquito larvae even in the presence of other kinds of foods.

Another important criterion for larvivorous fishes is their appearance. They should not be colorful or attractive to be hunted by other species, and most importantly, they should not have any nutritional value for humans, so that people do not eliminate them by eating (13). It is of course certain that finding a type that has all of the above parameters is difficult, but possible. The classification of

larval fishes is usually based on their feeding from the larval habitat. Some fishes, such as *Aplocheilus* and *Gambusia*, feed on the surface of water, and some of them such as *Amblypharyngodon mola*, *Danio* and *Rasbora*, feed on lower levels of water and some also feed on water depths, such as *Puntius* and *Colisa*. In the eradication program of malaria in Iran, the use of larvivorous fish varieties, especially *Gambusia*, was introduced as an auxiliary strategy along with other control methods of mosquito larvae (14). As in 1969, more than 1.5 million of the fish types were distributed in more than 3000 permanent water collections in Kermanshah (15). For example, after the release of fish in mosquito breeding sites, no population of larvae of *Anopheles* mosquitoes was found in stagnant waters around Kermanshah (15). Similarly, there was a significant decrease in larvae in Fars Province. In the village of Bahram Abad, Bushehr, in larval habitats where fish varieties were present, the larvae were not found, but there were numerous larvae in places where no larvivorous fish was present (15).


Therefore, based on the importance of the issue, the present study was a brief report of various types of larvivorous fish as a biological controller of mosquitoes in Iran and the world, which can provide interactions with prey/predator systems in aquatic ecosystems.

2. Materials and Methods

In this study, which was conducted as a non-systematic narrative review, keywords such as larvae, larvivorous fish, *Gambusia*, mosquito larvae and mosquito biological control were used. This study included an advanced searching in international scientific database including, Pub Med, Web of Science, Google

Scholar, Scopus, Elsevier, WHO and other internal scientific database including, Barakatks, Sicientific Information Database (SID), Medical Libraries (Medlib), Magiran Database, civilizational reference. A total of 9 scientific sources were collected, including books, articles, theses and reports that were then translated into Persian and English in the period from

1937 to 2017. Unrelated sources and articles were deleted and the resources related to our review were studied. Finally, 55 articles and the scientific sources were selected by following flowchart (Figure 1). Then, they were analyzed according to the purpose of the study.



3. Result

Table 1. Introduction, function and distribution of important species of larvivorous fish based on ecological characteristics and target mosquitoes

Fish species	Dispersion	ecology	Field tests	Distribution in Iran	Mosquito species
<i>Aphanius dispar</i>	India-Pakistan-Sind-Ethiopia-Palestine-Stretches of the Red Sea coast	Salt and sweet water, sewage and contaminated water, stagnant water, sewage wells.	About 3 fish per square meter of surface water eliminates mosquito larvae (19) 97% reduction of <i>Anopheles arabiensis</i> and <i>An.gambiae</i> (20) Removing <i>An.culicifacies adanensis</i> in Ethiopian Urban Area Wells (21)	Balochistan-Hormozgan (22)	<i>Cx. quinquefasciatus</i> <i>An. arabiensis</i> <i>An. gambiae complex</i> <i>An. culicifacie</i> <i>An. stephensi</i> <i>Aedes albopictus</i>
<i>Aplocheilus blockii</i>	East Coast of India, Sri Lanka	Fresh water, dwelling, small streams and rivers with poor plant growth, reservoirs, pools, foam and rivers, wells and wetlands	Reduced populations of <i>An.stephensi</i> larvae to 75% in the coastal belt (23) Control of <i>Aedes albopictus skuse</i> , resulting in control of chikonugonia fever in tankers, large reservoirs and large barrels of water <i>Ae. aegypti</i> Linnaeus, (Dengue Fever) (19)	was observed	<i>Ae. aegypti</i> <i>An. stephensi</i> <i>Aedes albopictus</i>
<i>Aplocheilus lineatus</i> <i>Aplocheilus panchax</i>	Indian, Peninsula and Sri Lanka India and Bengal, Orissa, Assam, Punjab, Sri Lanka, Malaya, Myanmar, Thailand and Indonesia.	Rice fields, aerial tanks, artificial vessels, reservoirs of pools, floor of rivers, marshes Quite durable and hardy and active Shallow water, sweet and salty water Wells, marshes, wetlands and contaminated drainage water, stagnant waters containing organic matter contamination	<i>An. culicifacies</i> control in irrigation channels, slow currents and low vegetation, river bed, rice fields <i>An. sundaicus</i> control in saline waters with algae, rice fields, reservoirs, wetlands, ponds, Control of <i>Cx. quinquefasciatus</i> in sewers, reservoirs, ponds, contaminated streams, <i>Cx. vishnui</i> control in rice fields, marshes, ponds, pools, streams, irrigation canals,	was observed was observed	<i>An. stephensi</i>
<i>Colisa fasciatus</i>	Myanmar, Punjab, Pakistan and across the peninsula of India	Rivers, lakes and rivers of fresh water Typically, the environment is covered with weeds	In the wells, complete removal of the larvae in one to two weeks after the arrival of 100 fish of the larvae (24)	was observed	Carriers of filariasis and malaria
<i>Colisa lalia</i>	North India	Slow flows of rivers, lakes, reservoirs Rivers and pools with appropriate vegetation	Production habitats and carriers such as stagnant weeds, margins of pools, tanks, holes and rice fields	Available (25)	Malaria Carriers
<i>Colisa sota</i>	North India	Rivers, pools and among plants Muddy water, stagnant and fresh water	Reduces larvae of mosquitoes in weed water, stagnant waters, and margins of pools, reservoirs, dead rivers, pits and rice fields.	was observed	Malaria Carriers
<i>Chanda nama</i>	India, Bangladesh and Burma (now Myanmar)	In freshly ground fresh vegetation Ponds, streams, streams, ponds	Suitable for controlling the variety of anopheles in sweet waters to control malaria	was observed	Malaria Carriers
<i>Oryzias melastigma</i>	India	Freshwater and coastal waters, pools, pools, rivers, canals and rivers, rice fields	Control of <i>Cx. vishnui</i> in rice fields, wetlands, ponds, pools, irrigation canals (24) <i>Anopheles</i> sp. and <i>Culex</i> sp. In rice fields, the decline in population on the sixth day, to 76.2%, in the subsequent days decreased from 98.3 to 100%, and from the 12th day on, decreased by 100%.	was observed	<i>An. culicifacies s.l.</i> <i>Cx. tritaeniorhynchus</i>
<i>Danio rerio</i>	North India, Bangladesh, Myanmar (Burma).	Slow flow, low-pitched wells, grass-clear transparent water,	In the rice farms, rich of <i>Anopheles</i> sp. and <i>Culex</i> sp. , On the sixth day, the reduction of larvae of age III, IV and IV by 86.8%, on the following days decreased from 92.4 to 99.3%, from the 12th day, decreased by 100% (24)	Available (Sanandaj-Golestan) (26)	<i>An. culicifacies s.l.</i> <i>Cx. tritaeniorhynchus</i>

<i>Macropodus cupanus</i>	East India, Sri Lanka, West Myanmar, Peninsula and Sumatra	In saline waters, contaminated canals, wetlands, shoals and shallow water, stagnant water, low food resistance or even lack of oxygen (27)	High level of hunting: Average hunting rate: 292.2 larvae per day	was not observed	<i>Culex quinquefasciatus</i>
<i>Carassius auratus (goldfish)</i>	China, Korea, Taiwan, Japan, Europe, Siberia, East Asia, India	Aquarium and decorative ponds	The number of larvae of <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i> and <i>Ar. subalbatus</i> was 193,188 and 132 per day (28) Effectively, <i>An. subpictus</i> larvae in reservoirs reduced density from 34.5 to 99.8% (28)	Available (25)	<i>An. subpictus</i>
<i>Gambusia affinis</i>	Native US Coastal Waters from southern New Jersey, Italy, Thailand, India	Fresh water, salt water Feeding aquatic and terrestrial insects (fish preference to terrestrial insects)	Daily feeding were 48, 51 and 31 larvae of <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i> and <i>Ar. subalbatus</i> (29) Mean hunting for each <i>Gambusia</i> fish is 7.65 <i>An. subpictus</i> larvae per day (30) Succeeding in the fight against malaria (15) Decreased <i>An. subpictus</i> to 96% (areas where fish have not been introduced, mosquito incidence increased from 55 to 92% (31) The best predator for <i>An. stephensi</i> larvae in air tanks (32) Introductions in rice fields have led to a sharp reduction in the density of <i>Anopheles</i> larvae (33)	Available (16)	<i>An. freeborni</i> <i>An. Pulcherrimus</i> <i>An. Stephensi</i> <i>An. Superpictus</i> <i>Cx. tritaeniorhynchus</i>
<i>Poecilia reticulata (guppy)</i>	America, India, West of Venezuela, Guyana	Low temperature intolerance Resistant to contamination	Feeding 18 to 32 <i>An. subpictus</i> larvae within 24 hours (34) Reduction of <i>An. aconitus</i> mosquito production in rice fields (35) <i>An. gambiae</i> larvae control in wipes, using 3-5 fishes up to 85% (36)	Available (25)	<i>An. gambiae s.s.</i> <i>An. stephensi</i> <i>An. subpictus</i>
<i>Nothobranchius guentheri</i>	East Africa, Mambaas Pengani River in Tanzania	Fish growth very fast About 20-100 eggs a day	The most suitable fish for malaria control using the Panama Canal (37)	was not observed	<i>Cx. quinquefasciatus</i>
<i>Xenentodon cancila</i>	Pakistan, India, Bangladesh, Sri Lanka, Myanmar and Thailand	In clear streams and permanent streams	Control of the fourth instar larvae <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i> and <i>Ar. subalbatus</i> , consumption of an average of 31, 28, 21 <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i> and <i>Ar. subalbatus</i> during the 24-hour period (29)	was not observed	<i>An. subpictus</i> <i>Cx. quinquefasciatus</i> <i>Ar. subalbatus</i>
<i>Oreochromis mossambica</i>	East Africa, India, Pakistan, Sri Lanka and so on	Rapid growth in saline water The temperature of the die is less than 10° Celsius	third and fourth stage of larvae <i>Cx. quinquefasciatus</i> and <i>An. culicifacies</i> with a rate of release of 5 fish per square meter (38)	Available (39)	<i>Cx. quinquefasciatus</i> <i>An. culicifacies</i>
<i>Oreochromis niloticus niloticus</i>	East Africa, West Africa, Nile River, Malaysia	Low salt tolerance (below c12) Suitable for agriculture in the tropical region	Introducing 20 fish in the field, reducing larval after 30 and 45 days, respectively. The removal of fish increased the larval congestion after 30 and 45 days (40)	was not observed	<i>An. gambiae</i> <i>An. Funestus</i> <i>An. stephensi</i>

Table 2. Examples of the use of a combination of two or more species of larvivorous fishes in the control of mosquito larvae

Species of larvivorous fish	Results in control of mosquito larvae
<i>Moroco oxycephalus</i> + <i>Misgurnus anguillicaudatus</i>	Control of <i>Anopheles sinensis</i> (41)
<i>Melanotaenia duboulayi</i> , <i>Retropinna semoni</i> , <i>Pseudomugil signifer</i>	Control of <i>Cx. annulirostris</i> , most used by <i>M. dbulaii</i> (42)
<i>Hypseleotris galii</i> , <i>Hypseleotris compressa</i> <i>Ambassis marianus</i> , <i>Gambusia holbrooki</i> <i>Pseudotropheus</i> , <i>Betta splendens</i> , <i>Osphronemus goramy</i> , <i>Pterophyllum scalare</i>	Control <i>Anopheles sinensis</i> Ranking: <i>P. tropheops</i> > <i>B. splendens</i> > <i>O. gorami</i> > <i>P. scalarae</i> Average daily feeding of larvae in 1 liter of water: 247, 238, 180 and 40 larva respectively (43)
<i>P. reticulata</i> - <i>Ga. affinis</i>	10 fish of <i>P. reticulata</i> and 5 fish of <i>Ga. Affinis</i> for feeding larvae of <i>Anopheles</i> , <i>Culex</i> and <i>Aedes</i> (40)
<i>G. affinis</i> - <i>Lebistes reticulatus</i>	Feeding on larva <i>An. subpictus</i> , the mean of feeding of larvae by <i>G. affinis</i> and <i>L. reticulatus</i> is 48 and 32, respectively (40)
<i>C. carpio</i> , <i>Ctenopharyngodon idella</i> , <i>Oreochromis niloticus</i> , <i>Clarias gariepinus</i>	Significant decrease in the fourth instar larvae <i>An. stephensi</i> in the 30 and 45 days after the arrival of the fish into the farm (44)
<i>Pseudomugil signifer</i> - <i>Gambusia holbrooki</i> <i>Aplocheilus lineatus</i> - <i>Oreochromis mossambicus</i>	Nutrition from the age of four larvae <i>Cx. annulrostris</i> (45) A reduction of 80-82% in the density of the <i>An. sinensis</i> larvae from the fifth week in the rice field (46)
<i>G. affinis</i> - <i>Tilapia nilotica nilotica</i> <i>Cyprinus carpio</i> - <i>Ctenopharyngodon idella</i> - <i>Tilapia</i>	Larva Control <i>An. freeborni</i> and <i>An. franciscanus</i> in the rice field (47) A reduction in <i>An. sinensis</i> during 150 to 170 days in rice fields and also reduction in the transmission of malaria (48)
<i>Carassius auratus</i> - <i>C. carpio</i> - <i>G. affinis</i> <i>G. affinis</i> , <i>Lethrinus reticulatus</i>	The effectiveness of mosquito control in rice farms in northern Italy (49) Larvae density of <i>An. Subpictus</i> in the presence of <i>G. affinis</i> and <i>L. reticulatus</i> was 25.7 to 0.36 and 23.7 to 0.5 (34)
<i>Lepisosteus tropicus</i> - <i>Astyanax fasciatus</i> - <i>Brycon guatemalensis</i> - <i>Ictalurus meridionalis</i> - <i>Poecilia reticulata</i> <i>Cnesterodon decemmaculatus</i> - <i>Jenynsia multidentata</i> <i>P. reticulata</i> - <i>G. affinis</i>	Control of <i>Aedes aegypti</i> larvae in water storage tanks (50) Control of fourth age of <i>Culex pipiens</i> (51) Control of <i>Cx. quinquefasciatus</i> larvae in drainage for about 3 months; reduction of larval congestion from 145 in each batch to 20 in each batch Control of 100% <i>An. culicifacies</i> and <i>An. stephensi</i> larvae (24)

4. Discussion

The use of biological control methods is expected to play an important role in future carrier management strategies. Unlike chemical pesticides, mosquito control results are unpredictable using biological agents. Therefore, developing and acquiring the necessary skills in this field is very necessary. Targeting interventions is essentially the most effective way to control vector-borne diseases, as most of the current efforts to develop new control tools are limited to laboratory scale (52).

In the application of larvivorous fish to control mosquitoes, two factors of fish adaptation and being native are very important. The most important adaptation of larvivorous fish in larval habitats is the ability to reproduce fish and feed the mosquito larvae. There are several factors that can influence the level of adaptation of larvivorous fish. For example, the use of a variety of pesticides and fertilizers can be harmful to fish used in irrigation fields (17). Aquatic plants, in turn, can also be effective in feeding fish, because it can hide mosquito larvae. Therefore, it is suggested that, before the release of larvivorous fish, an assessment of the effectiveness of one or more species of fish (especially native) is initially made (18). In this regard, fish behavior, the existence of minimum competition for food, proper water temperature, changes in water flow, suitable habitat for spawning, factors affecting species growth, such as diet, number of progenies per year, growth rate, body size, life span, and a lot of food have been introduced as the main factors promoting fish species creation.

On the other hand, it is important to note that introducing and releasing larvivorous fish as non-native species in a new ecosystem may have ecological hazards.

Because accidental releases or deliberate releases of fishes may cause some animal species to fall out of their natural range. Also, unplanned releases of larvivorous fish can cause unwanted problems with native counterparts and consequently reduce the efficiency of the mosquito control programs. Therefore, the introduction of a new fish strategy can create a lot of ecosystem hazards. For example, the species *Pseudorasbora parva* is highly competing with other fishes for food, and is a slaughterer of eggs and young fishes (16). Also, *Gambusia* fish can be negatively affected when it is not planted in waters beyond its native range, as it is greedy and highly invasive, with native fish for food and space life compete. It also feeds algae, zooplankton, aquatic insects, as well as eggs, young fish, amphibians, and even diatoms (53). On the other hand, *Gambusia* indirectly causes environmental changes as it increases the pH of the phytoplankton, increases water temperature and dissolved organic phosphorus, and decreases water clarity (54).

In the mosquito control program using larvivorous fish, it should be determined whether larvivorous fish have the ability to live even in drinking water and other waters. The residents are also trained so as not to accidentally kill the fish. When fish are used in different places for controlling mosquitos larvae, periodic addition of fish is essential to maintain the population needed to suppress larval mosquitoes. At times, in order to help the effectiveness and efficiency of larvivorous fish, the need for another larval control agent may also be felt. For example, simultaneous use of larvivorous fish with one of the commercial products of *Bacillus* bacteria is very promising to control mosquitoes in sites

where the fish cannot grow and reproduce sufficiently (55).

Considering that there are potential vectors of malaria in many regions of Iran and the climate of Iran has the potential to transmit the disease, it is suggested that, according to various ecological ecosystems of Iran, larvivorous fish that accept the environmental conditions of the area be used. In this regard, according to available findings, *Aphanius dispar* and *Danio rerio* can be used to control *Anopheles culicifacies* in areas, such as Sistan, South of Kerman, and Hormozgan. Also, *Gambusia affinis*, *Aplocheilus blochii*, *Aplocheilus panchax*, *Reiculate* (guppy) *Poecilia*, *Oreochromis niloticus* can be used to control *Anopheles stephensi* in southeast Iran. Finally, *Gambusia affinis* can be used to control *Anopheles superpictus* in the central plateau, mountainous regions of northern Iran and southern provinces of Iran.

Acknowledgments

The project was supported by a grant from Vice Chancellor for Research, Mazandaran University of Medical Sciences under grant number 3207. The researchers would like to express their gratitude to Vice Chancellor for Research, Mazandaran University of Medical Sciences for their financial support.

Conflict of interest

The authors also declare that they have no conflict of interest

References

1. Leta S, Beyene TJ, De Clercq EM, Amenu K, Kraemer MUG, Revie CW. Global risk mapping for major diseases transmitted by *Aedes aegypti* and *Aedes albopictus*. International journal of infectious diseases : IJID : official publication of the International Society for Infectious

- Diseases. 2018; 67: 25-35. PMID: 29196275
2. Benelli G, Jeffries CL, Walker T. Biological Control of Mosquito Vectors: Past, Present, and Future. Insects. 2016; 7(4): 1-18. PMID: 27706105
3. Collins LE, Blackwell A. The biology of Toxorhynchites mosquitoes and their potential as biocontrol agents. Biocontrol News and Information. 2000; 21(4): 105-16.
4. Milam C, Farris J, Wilhide J. Evaluating mosquito control pesticides for effect on target and nontarget organisms. Archives of Environmental Contamination and Toxicology. 2000; 39(3): 324-8. PMID: 10948282
5. Gratz N, Pal R. Malaria vector control: larviciding. Malaria: Principles and Practices of Malariology. 1988: 1213-26.
6. Mulla MS. Mosquito control investigations with emphasis on the integration of chemical and biological control in mosquito abatement. California Mosquito Control Association; 1961.
7. Raghavendra K, Subbarao S. Chemical insecticides in malaria vector control in India. ICMR bulletin. 2002; 32(10): 1-7.
8. Haas R, R. P. Mosquito larvivorous fishes. Bulletin of the ESA. 1984; 30(1): 17-25. DOI: <https://doi.org/10.1093/besa/30.1.17>.
9. Walton WE. Larvivorous fish including *Gambusia*. Journal of the American Mosquito Control Association. 2007; 23(2): 184-221. PMID: 17853606
10. Gerberich JB. Update of annotated bibliography of papers relating to the control of mosquitos by the use of fish for the years 1965-1981. Geneva: World Health Organization; 1985.
11. Waage J, Greathead D. Biological control: challenges and opportunities. Philosophical Transactions of the Royal Society of London B, Biological Sciences. 1988; 318(1189):111-28. DOI:<https://doi.org/10.1098/rstb.1988.0001>
12. Arthington A, Marshall C. Diet of the exotic mosquitofish, *Gambusia holbrooki*, in an Australian lake and potential for competition with indigenous fish species. Asian Fisheries Science. 1999; 12(1):1-16.
13. Job T. An investigation of the nutrition of the perches of the Madras coast. Records of the Indian Museum. 1940; 42:289-364.
14. Hora S, Mukherjee D. The absence of millions, *Lebistes reticulatus* (Peters) in

- India. Malaria Bureau No. 4. Health Bulletin No. 1938; 12:1-49.
15. Tabibzadeh I, Behbehani G, Nakhai R. Use of *Gambusia* fish in the malaria eradication programme of Iran. Bulletin of the World Health Organization. 1970; 43(4): 623. PMID: 5313079
 16. Esmaeili HR, Teimori A, Feridon O, Abbasi K, Brian WC. Alien and invasive freshwater fish species in Iran: Diversity, environmental impacts and management. Iranian Journal of Ichthyology. 2015; 1(2): 61-72. DOI: <http://dx.doi.org/10.22034/iji.v1i2.4>.
 17. Lacey LA, Lacey CM. The medical importance of riceland mosquitoes and their control using alternatives to chemical insecticides. Journal of the American Mosquito Control Association Supplement. 1990; 2:1-93. PMID: 1973949
 18. Blaustein L, Chase JM. Interactions between mosquito larvae and species that share the same trophic level. Annual review of entomology. 2007; 52(1): 489-507. PMID: 16978142
 19. Aatur-Rahim M. Observations on *Aphanius dispar* (Ruppell, 1828), a mosquito larvivorous fish in Riyadh, Saudi Arabia. Annals of tropical medicine and parasitology. 1981; 75(3): 359-62. PMID: 6895451
 20. Louis JP, Albert JP. Malaria in the Republic of Djibouti. Strategy for control using a biological antilarval campaign: indigenous larvivorous fishes (*Aphanius dispar*) and bacterial toxins. Medecine tropicale : revue du Corps de sante colonial. 1988; 48(2): 127-31. PMID: 3043137
 21. Fletcher M, Teklehaimanot A, Yemane G. Control of mosquito larvae in the port city of Assab by an indigenous larvivorous fish, *Aphanius dispar*. Acta tropica. 1992; 52(2-3): 155-66. PMID: 1363180
 22. Reichenbacher B, Kamrani E, Esmaeili HR, Teimori A. The endangered cyprinodont *Aphanius ginaonis* (Holly, 1929) from southern Iran is a valid species: evidence from otolith morphology. Environmental Biology of Fishes. 2009; 86(4): 507.
 23. Kumar A, Sharma VP, Sumodan PK, Thavaselvam D. Field trials of biolarvicide *Bacillus thuringiensis* var. *israelensis* strain 164 and the larvivorous fish *Aplocheilus blocki* against *Anopheles stephensi* for malaria control in Goa, India. Journal of the American Mosquito Control Association. 1998; 14(4): 457-62. PMID: 10084141
 24. Sharma V, Ghosh A, editors. Larvivorous fishes of inland ecosystems. Proceedings of the MRC-CIFRI Workshop Delhi: India Malaria Research Centre (ICMR) publication; 1994.
 25. Mood S, Ebrahimzadeh Mousavi H, Mokhayer B, Ahmadi M, Soltani M, Sharifpour I. *Centrocestus formosanus* metacercarial infection of four ornamental fish species imported into Iran. Bulletin of the European Association of Fish Pathologists. 2010; 30(30): 146-9.
 26. Mansouri B, Maleki A, Johari SA, Reshahmanish N. Effects of cobalt oxide nanoparticles and cobalt ions on gill histopathology of zebrafish (*Danio rerio*). Aquaculture, Aquarium, Conservation & Legislation. International Journal of the Bioflux Society (AAFL Bioflux). 2015; 8(3): 438-444.
 27. Mathayan S, Muthukrishnan J, Heleenal G. Studies on predation on mosquito larvae by the fish *Macropodus cupanus*. Hydrobiologia. 1980; 75(3): 255-8. DOI: <https://doi.org/10.1007/BF00006490>.
 28. Chatterjee S, Das S, Chandra G. Gold fish (*Carrasius auratus*) as a strong larval predator of mosquito. Transactions of the Zoological Society of London. 1997; 1: 112-4.
 29. Chatterjee S, Chandra G. Laboratory trials on the feeding pattern of *Anopheles subpictus*, *Culex quinquefasciatus* and *Armigeres subalbatus* larvae by *Gambusia affinis*. Science and Culture. 1997; 63: 51-2.
 30. Menon PK, Rajagopalan PK. Control of mosquito breeding in wells by using *Gambusia affinis* and *Aplocheilus blochii* in Pondicherry town. The Indian journal of medical research. 1978; 68: 927-33. PMID: 582034
 31. Bheema Rao US, Krishnamoorthy K, Reddy CB, Panicker KN. Feasibility of mosquito larval control in casuarina pits using *Gambusia affinis*. The Indian journal of medical research. 1982; 76: 684-8. PMID: 6897726
 32. Kant R, Pandey SD, Sharma SK. Role of biological agents for the control of mosquito breeding in rice fields. Indian journal of malariology. 1996; 33(4): 209-15. PMID: 9125835

33. Rafatjah H, Arata A. The use of larvivorous fish in antimalaria programmes. Geneva. World Health Organization, (unpublished document MAL/WP/756 Rev 1). 1975.
34. Chatterjee S, Chandra G. Feeding pattern on *Gambusia affinis* and *Lebistes reticulatus* on *Anopheles subpictus* larvae in the laboratory and field condition. *Journal of applied zoological researches*. 1997; 8(2): 152-3.
35. Nalim S, Boewono D, Haliman A, Winoto E. Control demonstration of the rice field breeding mosquito *Anopheles aconitus* Donitz in Central Java (Indonesia), using *Poecilia reticulata* through community participation, 1: Experimental design and concept. *Buletin Penelitian Kesehatan (Indonesia)*. 1985.
36. Sabatinelli G, Blanchy S, Majori G, Papakay M. Impact of the use of larvivorous fish *Poecilia reticulata* on the transmission of malaria in FIR of Comoros. *Annales de parasitologie humaine et comparee*. 1991; 66(2): 84-8. PMID: 1952700
37. Vanderplank F. *Nothobranchius* and *Barbus* Species: indigenous anti-malarial Fish in East Africa. *East African Medical Journal*. 1941; 17(10): 431-436.
38. Vaidyanathan R, Feldlaufer MF. Bed bug detection: current technologies and future directions. *The American journal of tropical medicine and hygiene*. 2013; 88(4): 619-25. PMID: 23553226
39. Marjani M, Jamili S, Mostafavi P, Ramin M, A. M. Influence of 17-alpha methyl testosterone on masculinization and growth in tilapia (*Oreochromis mossambicus*). *Journal of fisheries and aquatic science*. 2009; 4(1): 71-4. DOI: 10.3923/jfas.2009.71.74.
40. Ghosh A, Bhattacharjee I, G. C. Biocontrol of larval mosquitoes by *Oreochromis niloticus niloticus*. *Journal of Applied Zoological Researches*. 2006; 17(1): 114-6.
41. Kim HC, Kim MS, HS Y. Biological control of vector mosquitoes by the use of fish predators, *Moroco oxycephalus* and *Misgurnus anguillicaudatus* in the laboratory and semi-field rice paddy. *Korean Journal of Entomology*. 1994.
42. Hurst TP, Brown MD, Kay BH. Laboratory evaluation of the predation efficacy of native Australian fish on *Culex annulirostris* (Diptera: Culicidae). *Journal of the American Mosquito Control Association*. 2004; 20(3): 286-91. PMID: 15532929
43. Ghosh A, Bhattacharjee I, Ganguly M, Mondal S, G. C. Efficacy of some common aquarium fishes as biocontrol agent of preadult mosquitoes. *Buletin Penelitian Kesehatan (Indonesia)*. 2004; 32(4): 144-9.
44. Ghosh A, MANDAL S, Bhattacharjee I, G. C. Biological control of vector mosquitoes by some common exotic fish predators. *Turkish journal of biology*. 2005; 29(3): 167-71.
45. Willems KJ, Webb CE, Russell RC. A comparison of mosquito predation by the fish *Pseudomugil signifier* Kner and *Gambusia holbrooki* (Girard) in laboratory trials. *Journal of vector ecology: journal of the Society for Vector Ecology*. 2005; 30(1): 87-90. PMID: 16007960
46. Yu HS, JH. L. Biological control of malaria vector (*Anopheles sinensis* Wied.) by combined use of larvivorous fish (*Aplocheilus latipes*) and herbivorous hybrid (*Tilapia mossambicus niloticus*) in rice paddies of Korea. *Korean journal of applied entomology*. 1989; 28(4): 229-36.
47. Kramer V. The ecology and biological control of mosquitoes [*Culex tarsalis*, *Anopheles freeborni* and *A. franciscanus* [*A. pseudopunctipennis franciscanus*]] in California wild and white rice fields [using *Bacillus thuringiensis* and larvivorous fish]. *Dissertation Abstracts International B, Sciences and Engineering*. 1989;49.
48. Wu N, Liao GH, Li DF, Luo YL, Zhong GM. The advantages of mosquito biocontrol by stocking edible fish in rice paddies. *The Southeast Asian journal of tropical medicine and public health*. 1991; 22(3): 436-42. PMID: 1818398
49. Bellini R, Veronesi R, M. R. Efficacy of various fish species (*Carassius auratus* [L.], *Cyprinus carpio* [L.], *Gambusia affinis* [Baird and Girard]) in the control of rice field mosquitoes in Northern Italy. *Bulletin of the Society for Vector Ecology*. 1994; 19: 87-99.
50. Martinez-Ibarra J, Guillén YG, Arredondo-Jimenez J, M. R-L. Indigenous fish species for the control of *Aedes aegypti* in water storage tanks in Southern Mexico. *BioControl*. 2002; 47(4): 481-6. DOI: <https://doi.org/10.1023/A:101569183>.
51. Marti GA, Azpelicueta Mde L, Tranchida MC, Pelizza SA, Garcia JJ. Predation

- efficiency of indigenous larvivorous fish species on *Culex pipiens* L. larvae (Diptera: Culicidae) in drainage ditches in Argentina. *Journal of vector ecology : journal of the Society for Vector Ecology*. 2006; 31(1): 102-6. PMID: 16859096
52. Haq S, Yadav R, V. K. Developing larvivorous fish network for mosquito control in urban areas: A case study. *ICMR Bulletin*. 2003; 33(7): 69-73.
53. García- Berthou E. Food of introduced mosquitofish: ontogenetic diet shift and prey selection. *Journal of Fish Biology*. 1999; 55(1): 135-47. DOI: <https://doi.org/10.1111/j.1095-8649.1999.tb00663.x>.
54. Hurlbert SH, Zedler J, Fairbanks D. Ecosystem Alteration by Mosquitofish (*Gambusia affinis*) Predation. *Science*. 1972;175(4022):639-41. DOI: 10.1126/science.175.4022.639.
55. Walker K, Lynch M. Contributions of Anopheles larval control to malaria suppression in tropical Africa: review of achievements and potential. *Medical and veterinary entomology*. 2007; 21(1): 2-21. DOI: 10.1111/j.1365-2915.2007.00674.x.